

National Immunisation Program

July 2020 South Australia Schedule[#]

† CAUTION brand name similarity
 ■ Additional vaccines

CHILDREN < 5 YEARS OF AGE

Age	Antigen	Vaccine Brand	Route	Important Information
Birth	Hepatitis B	Engerix B [®] paediatric OR HB Vax II [®] paediatric	IM IM	Within 7 days of birth (ideally within 24 hours) No catch-up required
6 weeks	DTPa-hepB-IPV-Hib 13vPCV Rotavirus Meningococcal B	Infanrix [®] hexa [†] Prevenar 13 [®] Rotarix [®] Bexsero [®]	IM IM Oral IM	Rotarix – dose 1 limited to 6-14 weeks of age Bexsero – prophylactic paracetamol required
MAR*	Meningococcal ACWY vaccine is recommended for children with specific MAR conditions. Refer to the online Australian Immunisation Handbook.			
4 months	DTPa-hepB-IPV-Hib 13vPCV Rotavirus Meningococcal B	Infanrix [®] hexa [†] Prevenar 13 [®] Rotarix [®] Bexsero [®]	IM IM Oral IM	Rotarix – dose 2 limited to 10-24 weeks of age Bexsero – prophylactic paracetamol required
MAR	Meningococcal ACWY vaccine is recommended for children with specific MAR conditions. Refer to the online Australian Immunisation Handbook.			
6 months	DTPa-hepB-IPV-Hib	Infanrix [®] hexa [†]	IM	
Aboriginal**	13vPCV	Prevenar 13 [®]	IM	
MAR	Pneumococcal, meningococcal ACWY and/or meningococcal B vaccines are recommended for children with specific MAR conditions. Refer to the online Australian Immunisation Handbook.			
12 months	4vMenCV (ACWY) MMR 13vPCV Meningococcal B	Nimenrix [®] MMR II [®] OR Priorix [®] Prevenar 13 [®] Bexsero [®]	IM Subcut/IM IM IM	Bexsero – prophylactic paracetamol required
MAR	Hepatitis B vaccine is recommended for children who were low birth weight (<2000g) and/or preterm (<32 weeks)			
18 months	Hib DTPa MMRV	Act-HIB [®] Tripacel [®] OR Infanrix [†] Priorix Tetra [®] OR ProQuad [®]	IM IM Subcut/IM Subcut/IM	
Aboriginal	Hepatitis A	Vaqa [®] paediatric	IM	
4 years	DTPa/IPV	Infanrix IPV [®] OR Quadracel [®]	IM IM	
Aboriginal	Hepatitis A 23vPPV	Vaqa [®] paediatric Pneumovax 23 [®]	IM IM	
MAR	Pneumococcal vaccine is recommended for children with specific MAR conditions. Refer to the online Australian Immunisation Handbook.			

Additional *Haemophilus Influenzae Type B (Hib)*, meningococcal and/or pneumococcal vaccines are funded for specific MAR groups. The recommendations are very specific. Refer to the online Australian Immunisation Handbook for details and further information.

CHILDREN ≥ 5 YEARS OF AGE / ADOLESCENTS / ADULTS

Age	Antigen	Vaccine Brand	Route	Important Information
9 – 14 years Aboriginal	23vPPV	Pneumovax 23 [®]	IM	5 – 10 years after the previous dose of 23vPPV
Year 8 students	dTpa HPV	Boostrix [®] Gardasil [®] 9	IM IM	9 to <15 years, 2 doses HPV with a 6 month interval
Year 10 students	4vMenCV (ACWY) Meningococcal B	Nimenrix [®] Bexsero [®]	IM IM	2 doses Bexsero with a minimum 8 week interval
50 years Aboriginal	13vPCV	Prevenar 13 [®]	IM	
Aboriginal	23vPPV	Pneumovax 23 [®]	IM	First dose 23vPPV should be given 12 months after the 13vPCV Second dose of 23vPPV 5-10 years later No more than 2 doses of 23vPPV to be given in a lifetime
≥ 70 years	13vPCV VSZ (shingles)	Prevenar 13 [®] Zostavax [®]	IM Subcut	Minimum interval of 12 months after any previous dose of 23vPPV Catch-up program for aged 71-79 years until 31 October 2021
Pregnancy	dTpa	Adacel [®] OR Boostrix [®]	IM	Each pregnancy from 20 weeks gestation

[#] Refer to the Annual Funded Influenza Program South Australia Schedule for influenza vaccine information

* Medically at risk (MAR)

** The term Aboriginal is inclusive of Aboriginal and Torres Strait Islander people

For more information
 Immunisation Section, SA Health
sahealth.sa.gov.au/immunisationprovider

Public-I2-A1
 @Department for Health and Wellbeing,
 Government of South Australia. All rights reserved.
 Printed May 2020. FIS: 20063.1



<https://creativecommons.org/licenses>